



Online article and related content  
current as of July 19, 2010.

## National Survey of Internal Medicine Residency Program Directors Regarding Problem Residents

David C. Yao; Scott M. Wright

*JAMA*. 2000;284(9):1099-1104 (doi:10.1001/jama.284.9.1099)

<http://jama.ama-assn.org/cgi/content/full/284/9/1099>

Correction	<a href="#">Contact me if this article is corrected.</a>
Citations	<a href="#">This article has been cited 31 times.</a> <a href="#">Contact me when this article is cited.</a>
Topic collections	Public Health; Public Health, Other; Quality of Care; Quality of Care, Other <a href="#">Contact me when new articles are published in these topic areas.</a>
Related Articles published in the same issue	September 6, 2000 <a href="#">JAMA. 2000;284(9):1175.</a>
Related Letters	Alcohol and Substance Abuse Among Medical Residents <a href="#">J. Mitchell Simson et al. JAMA. 2000;284(22):2874.</a>

Subscribe  
<http://jama.com/subscribe>

Permissions  
[permissions@ama-assn.org](mailto:permissions@ama-assn.org)  
<http://pubs.ama-assn.org/misc/permissions.dtl>

Email Alerts  
<http://jamaarchives.com/alerts>

Reprints/E-prints  
[reprints@ama-assn.org](mailto:reprints@ama-assn.org)

# National Survey of Internal Medicine Residency Program Directors Regarding Problem Residents

David C. Yao, MD, MPH

Scott M. Wright, MD

SINCE 1972, THE AMERICAN Board of Internal Medicine (ABIM) has relied on residency programs to evaluate the readiness of eligible candidates for certification. Accordingly, attempts have been made to standardize resident evaluation at the program level, and to identify those residents who fail to meet the evaluation norms. The ABIM defines a *problem resident* as "a trainee who demonstrates a significant enough problem that requires intervention by someone of authority, usually the program director or chief resident."<sup>1</sup> The problem is often manifested in 1 or more of the ABIM's 7 areas that relate to clinical competency: clinical judgment, medical knowledge, clinical skills, humanistic qualities, professional attitudes and behavior, medical care, and moral and ethical behavior. Residency program directors predictably have to work with problem residents and must consider the negative impact they may have on the residency program.

Although problem residents pose challenges for residency programs, there is little information on their epidemiology or management. A number of articles have addressed specific issues such as impairment due to emotional stress and substance abuse.<sup>2-25</sup> To gain more understanding of the prevalence, identification, management, and prevention of problem residents, we con-

**Context** Internal medicine residency training is demanding and residents can experience a wide variety of professional and personal difficulties. A *problem resident* is defined by the American Board of Internal Medicine as "a trainee who demonstrates a significant enough problem that requires intervention by someone of authority." Data are sparse regarding identification and management of such residents.

**Objective** To gain more understanding of the prevalence, identification, management, and prevention of problem residents within US internal medicine residency programs.

**Design, Setting, and Participants** Mailed survey of all 404 internal medicine residency program directors in the United States in October 1999, of whom 298 (74%) responded.

**Main Outcome Measures** Prevalence of problem residents; type of problems encountered; factors associated with identification and management of problem residents.

**Results** The mean point prevalence of problem residents during academic year 1998-1999 was 6.9% (SD, 5.7%; range, 0%-39%), and 94% of programs had problem residents. The most frequently reported difficulties of problem residents were insufficient medical knowledge (48%), poor clinical judgment (44%), and inefficient use of time (44%). Stressors and depression were the most frequently identified underlying problems (42% and 24%, respectively). The most frequent processes by which problem residents were discovered included direct observation (82%) and critical incidents (59%). Chief residents and attending physicians most frequently identified problem residents (84% and 76%, respectively); problem residents rarely identified themselves (2%). Many program directors believed that residents who are from an underrepresented minority, are international medical graduates, or are older than 35 years are at increased risk of being identified as a problem resident ( $P < .05$ ). Program directors believed that frequent feedback sessions (65%) and an assigned mentor for structured supervision (53%) were the most helpful interventions.

**Conclusion** Nearly all internal medicine residency programs in this sample had problem residents, whose presenting characteristics and underlying issues were diverse and complex.

JAMA. 2000;284:1099-1104

www.jama.com

ducted a national survey of internal medicine residency program directors.

## METHODS

### Participants

Program directors of all internal medicine residency programs in the United

**Author Affiliations:** Division of General Internal Medicine, Johns Hopkins Bayview Medical Center, and the Johns Hopkins University School of Medicine, Baltimore, Md. Dr Yao is now with the Medical Education Program, University of California, San Francisco in Fresno. **Corresponding Author and Reprints:** David C. Yao, MD, MPH, UCSF-Fresno, Medical Education Program, University Medical Center, 445 S Cedar Ave, Fresno, CA 93702 (e-mail: dcyao@alum.mit.edu).

**Table 1.** Frequency of Apparent Deficiencies and Underlying Causes in Problem Residents

	Half of the Time or More Frequently, %
Apparent deficiencies	
Insufficient medical knowledge	48
Poor clinical judgment	44
Inefficient use of time	44
Inappropriate interaction with colleagues or staff	39
Provision of poor or inadequate medical care to patients	36
Unsatisfactory clinical skills	31
Unsatisfactory humanistic behavior with patients	23
Excessive and unexplained tardiness or absences	21
Unacceptable moral or ethical behaviors	15
Underlying causes	
Situational, personal, or professional stresses	42
Depression	24
Cognitive dysfunction	8
Other psychiatric illness	5
Substance abuse, other than alcohol	1
Alcohol abuse	<1

States and Puerto Rico were targeted for inclusion in this study. A list of the 404 accredited internal medicine programs was obtained from the American Medical Association's Graduate Medical Education Directory.<sup>26</sup>

### Survey Content

For this study, we developed a detailed 89-item questionnaire focusing on the identification, prevalence, management, and prevention of problem residents. Relevant issues pertaining to problem residents were identified by reviewing the published literature using MEDLINE, PsychINFO, ERIC, and HealthSTAR and by conducting focused interviews with 6 internal medicine residency program directors in Maryland.

The questionnaire was organized into 7 main sections and used the ABIM's definition of a "problem resident."<sup>1</sup> The first section addressed the problematic behaviors and deficiencies noted among problem residents and their underlying causes. The second section explored how and by whom these residents were identified. The third section collected data about individual residency programs and

the prevalence of problem residents within each program. The fourth section addressed the remedial actions used to improve the performance of problem residents. The fifth section addressed the disposition and outcome of problem residents. The sixth section asked about measures that program directors use to screen for potential problem residents during the residency application process. The final section collected data about each residency program and the program directors. The majority of the questionnaire items used a 7-point word-anchored Likert scale ("never, rarely, occasionally, half of the time, frequently, almost always, always").

### Survey Administration

After pilot testing the questionnaire with selected program directors and attending physicians, the confidential questionnaire was mailed to each program director in October 1999. Reminder postcards and e-mail messages were sent 4 weeks later, and a second mailing was sent to those who did not respond within 8 weeks. Finally, for those who still did not respond, follow-up telephone calls and faxes of the questionnaire were used to encourage full participation.

### Data Analysis

Survey responses were reviewed and entered into a computerized database.<sup>27</sup> The frequency distribution of responses was examined to find evidence of nonnormality, outliers, and other irregularities in the data. Descriptive statistics (mean, median, range, and SD) were used to summarize the responses to all questions. The sign test was used to compare various resident characteristics that may place an individual resident at increased risk of being identified as a problem resident. Spearman correlations were used to determine whether program characteristics (percentage of male residents, minorities, etc) influenced the risk of housestaff with the same characteristics being identified as problem residents.  $\chi^2$  Analyses were performed to detect differences in the characteristics, identification, and management of problem

residents in community vs university teaching hospitals. Residency programs at Veterans Affairs and military hospitals constituted 5% of the sample and were not included in this subset analysis.

### RESULTS

Of the 404 program directors in the mailing, 298 returned a completed questionnaire, for a 74% response rate. On average, program directors had been serving in that capacity for 7 (SD, 6) years. Eighty-four percent of respondents were male and 16% were female (vs 86% male, 14% female for nonrespondents;  $P = .49$ ). Fifty-eight percent of respondents were from community-based teaching hospitals, 37% from university-based hospitals, and 5% from Veterans Affairs and military teaching hospitals (the corresponding percentages for nonrespondents were 59%, 39%, and 2%;  $P = .30$ ).

The mean point prevalence of problem residents per program in the 1998-1999 academic year as reported by the program directors was 6.9% (SD, 5.7%; range, 0%-39%). Six percent of responding program directors stated that there were no problem residents in their programs. The majority (64%) stated that the prevalence had remained constant during their tenure as program directors. Twenty-one percent believed that the prevalence had decreased, which some respondents attributed to a better pool of applicants, an improved selection process, and the closer surveillance of residents with frequent feedback. Fifteen percent reported that the prevalence had increased; some of these respondents thought that this was due to a sense of entitlement by the recent cohort of residents and increased complexity of patient illness.

### Apparent and Underlying Causes

The most prevalent deficiencies identified by respondents were insufficient medical knowledge, poor clinical judgment, and inefficient use of time (TABLE 1). These were identified by 48%, 44%, and 44% of program direc-

tors, respectively, as being present “half of the time or more frequently” in problem residents. Situational, personal, or professional stresses and depression were believed to be the most frequently recognized underlying causes (Table 1).

**Identification**

Sixty-seven percent of program directors stated that the return rate of attending physicians’ evaluation of residents was more than 75% in their programs. While the majority of program directors (74%) did not think that low evaluation return rate had been an issue in identifying problem residents, 60% strongly agreed or agreed that it had been difficult to convince problem residents of their deficiencies because of a lack of honest and accurate written evaluations from attending physicians. TABLE 2 lists the processes and personnel through which problem residents were identified. Eighty-two percent of program directors reported that direct observation in clinical setting was the most frequent method by which problem residents were identified. Critical incidents and poor performance at morning report or other conferences also frequently led to the identification of problem residents. Chief residents, attending physicians, and program directors most frequently discovered problem residents, although other residents and nursing staff also did so. Problem residents seldom identified themselves as such (Table 2).

Program directors were asked about the influence of certain personal characteristics of residents on the likelihood that a trainee would be identified as a problem resident. Most program directors did not believe that any of these characteristics had a bearing on this matter. Although these variables were thought to be predictive by some program directors, the proportion of program directors who perceived a given factor as decreasing a resident’s risk of becoming a problem resident was always smaller than the proportion who perceived that factor

to increase the risk (TABLE 3). Residents who were older than 35 years, international medical graduates (IMGs), and those of an underrepresented minority (this term was not defined in the survey instrument) were considered to be at greatest risk of being identified as a problem resident. Based on the perception of the program directors, these 3 factors were significantly more likely to be associated with being identified as a problem resident ( $P < .05$  for each) compared with other characteristics, such as single marital status, male sex, or being a parent. However, the correlation test results showed that as the percentage of older residents, minorities, and IMGs increased within programs, program directors were less concerned about the potential for poor performance of these residents. Furthermore, IMGs were significantly less likely to be identified as problem residents in programs with high proportions of IMGs ( $P = .008$ ).

**Management**

Interventions that were reported by program directors as having helped to correct the identified problem, in situations in which the intervention was applicable, are listed in TABLE 4. The most frequent helpful intervention for problem residents was more frequent feedback sessions, followed by assigning a mentor for structured supervision, probation, psychiatric or psychological counseling, strict behavioral guidelines, and remedial didactic curricula.

When asked about their comfort level in dealing with problem residents, 88% of program directors were confident of their program’s ability to identify them. Seventy-six percent were confident in their program’s ability to diagnose or understand the causes of a problem resident’s difficulties, and 58% were confident of their program’s capability to manage problem residents.

In terms of outcomes seen in problem residents, program directors estimated that 57% finished their resi-

**Table 2.** Frequency of Processes and Individuals Who Identify Problem Residents

	Half of the Time or More Frequently, %
Processes that identified problem residents	
Direct observation in clinical settings	82
Critical incident	59
Poor performance at morning report/conferences	45
Neglecting patient care responsibility	33
Chart review/medical record audit	26
In-training examination	23
Mini-CEX (clinical evaluation exercise)	8
Individuals who identified problem residents	
Chief residents	84
Attending physicians, through verbal comments	76
Program director	74
Other residents	49
Attending physicians, through written evaluations	41
Nursing staff	31
Self	2
Patients and their families	2

**Table 3.** Perceived Effect of 6 Factors on the Chance That a Resident Will Be Identified as a Problem Resident\*

Factors	Risk, %		
	Decrease	No Effect	Increase
Underrepresented minority†	1	69	28
International medical graduate†	6	53	29
>35 Years†	5	59	33
Single	4	84	11
Male	2	82	16
Parent	3	76	19

\*Some percentages do not add to 100% because program directors could also choose “not applicable (no such residents).”  
 †Statistically significantly ( $P < .05$ ) increases the chance of being identified as a problem resident when compared to the bottom 3 variables.

dency training at the appropriate time, 18% continued in the residency program but required additional training time, 9% transferred to another internal medicine residency program, 10% transferred to a different residency pro-

gram (other than internal medicine), and 4% left the profession of medicine.

With respect to the legal repercussions of interacting with problem residents, 49% of program directors acknowledged threats of litigation as a result of their dealings with problem residents during their tenure as program directors. Fifteen percent of program directors indicated that their interactions with problem residents have resulted in actual lawsuits.

**Table 4.** Frequency With Which Selected Interventions Have Helped Problem Residents Correct Identified Problems

Interventions	Half of the Time or More Frequently, %
More frequent feedback sessions	65
Assigned mentor for structured supervision	53
Probation	35
Psychiatric/psychological counseling	35
Strict behavioral guidelines	32
Remedial didactic curriculum	28
Leave of absence	9
Formal psychomotor function testing/learning assessment	7
Substance abuse rehabilitation program	7

**Table 5.** Perceived Helpfulness of Selected Factors in Screening for Potential Problem Residents During Application

Factors	Half of the Time or More Frequently, %
"Audition" rotation while a medical student	38
Interviews	36
Board scores	33
School records	28
Dean's letter	20
Recommendation letters	19
Reputation of the medical school	16

**Table 6.** Differences in Problem Resident Identification and Management Between University- and Community-Based Teaching Hospitals\*

Characteristics	University, %	Community, %
Underlying cause		
Depression		
Half of the time or more	30	19
Identification		
Program directors		
Half of the time or more	63	80
Chief residents		
Half of the time or more	90	79
Mini-CEX		
Half of the time or more	4	16
Management		
Counseling		
Half of the time or more	45	30
Retreats for interns and residents	61	38
Advisory system	65	78
Prevention		
Board scores		
Half of the time or more	25	42

\*All factors statistically significantly different at  $P < .05$ . CEX indicates clinical evaluation exercise.

and enhance the development of problem residents included regular meetings with program director and/or chief residents (94%), timely evaluations from attending physicians (87%), regular meetings of attending physicians with program director and/or department chairperson before and after ward rotations (49%), and switching rotations if the problem resident is particularly stressed and unable to perform at a high level (29%).

Program directors were questioned about the use of multiple factors in screening for potential problem residents during the application process. "Audition" rotations, interviews, board scores, and school records (eg, grade point average, ranking) were reported most frequently as being helpful (TABLE 5).

**Subset Analyses**

Subset analyses were performed to compare university- and community-based residency programs. The prevalence of problem residents was higher in community-based programs (8.1%) than in university-based programs (5.0%) ( $P < .001$ ). Community-based residency programs had more residents who are male ( $P = .003$ ), underrepresented minorities ( $P = .002$ ), older than 35 years ( $P = .009$ ), IMGs ( $P < .001$ ), and parents ( $P < .001$ ). TABLE 6 shows additional differences between university- and community-based residency programs on issues related to problem residents. Program directors from university-based hospitals identified depression as being more common among their problem residents. Program directors at community-based programs reported that they more frequently identified problem residents, while those at university-based programs reported that chief residents more frequently identified problem residents. Psychiatric or psychological counseling was reported as more frequently helping problem residents in university-based programs, perhaps reflecting the higher frequency of reported depression in these programs. In screening applicants, community-based programs used board scores as a predictor

**Preventive Measures**

Program directors were asked to describe the measures in place that may benefit and support the well-being of their housestaff. More than 90% of the programs have the following: (1) orientation at the beginning of residency regarding expectations and supportive resources, (2) limited number of admissions on call days, (3) structured days off during ward rotations, (4) planned social events, and (5) parental leave policies. Seventy-three percent of programs have an advisory system, 48% sponsor retreats for interns and residents (ie, outings away from the hospital setting to assist in personal and professional development), and 35% offer intern/resident support groups or personal awareness groups.

Measures that program directors reported using to prevent deterioration

more frequently than did university-based programs.

## COMMENT

Problem residents are moderately prevalent among internal medicine residency training programs. However, heightened awareness by medical educators may have resulted in an even higher estimate of the prevalence of problem residents. This may explain why our estimate was lower than that of the ABIM, which through on-site visits had estimated that 8% to 15% of residents have serious problems.<sup>28</sup> Many of our findings, such as identification, apparent and underlying causes, management, and preventive measures, represent new data.

Chief residents and program directors most often identify problem residents. The finding that problem residents do not generally identify themselves raises the question of whether there is accurate self-evaluation among these house officers. Other factors that might play a role are denial, embarrassment, fear of consequences, or a concern about the inability to improve in a problem area.

Program directors acknowledged that multiple apparent deficiencies and underlying causes frequently contribute to the suboptimal performance of problem residents. While the apparent deficiencies manifested by the problem residents are often noted first, underlying issues may be at the root of the problem. Heightened sensitivity to situational, emotional, cognitive, and cultural difficulties may lead to improvements in the apparent deficiencies.

Management of problem residents can be difficult, and litigation is a possible consequence.<sup>29</sup> Both threatened and actual lawsuits were reported by a surprisingly high number of respondents. The ABIM emphasizes the importance of documentation and due process; many residency programs can likely improve in this area.<sup>30</sup>

Reliable predictors of individuals at risk for becoming problem residents could be used by programs in selecting candidates and targeting preven-

tive strategies. "Audition rotations," interviews, board scores, and school records were reported as useful processes in screening for problem residents. Unfortunately, there is a lack of information on the efficacy of these variables or processes in predicting which applicants will become problem residents. It may be difficult to predict individual responses to the stresses of residency, and perhaps more emphasis should be placed on detecting problem residents early.

The majority of the program directors believed that personal factors do not affect the chance that an individual house officer will be identified as a problem resident. Among the program directors who did not share this view, residents who were older, underrepresented minorities, or IMGs were thought to be at greatest risk for becoming problem residents. These results raise the issue of whether medical students or housestaff with these characteristics may be discriminated against during the application process or training. Cultural differences, language difficulties experienced by some IMGs, expectations, and previous experiences may place these groups at a disadvantage at the beginning of residency training. Discrimination has been described in residency training programs, and program directors need to be cognizant of these findings.<sup>31,32</sup>

Several limitations of this study should be considered. First, the results are derived from a cross-sectional survey of program directors that relied on self-report, and therefore, causal relationships and efficacy of management strategies could not be assessed. Second, since program directors were asked to think about their experiences with problem residents during their tenure, recall bias may have occurred as program directors may preferentially remember specific cases or residents who were most challenging. Third, because of the sensitive nature of the content addressed in this confidential study, some respondents may have been overly cautious. Fourth, there are some areas, such as the prevalence of underlying factors, in which it would have

been clearly preferable to ask the problem residents themselves. Finally, the results of this study may not be generalizable to training programs other than internal medicine.

Further research on problem residents may address issues such as the identification of reliable and valid predictors, trials of specific preventive and management strategies, and detailed assessments of the outcomes and disposition of problem residents. Investigations from the perspective of the problem residents themselves would be useful. Furthermore, among residency programs in which the prevalence of problem residents is significantly higher than the national mean, consideration of the system as the problem (as opposed to the residents) should be investigated.

In summary, our findings show that the vast majority of internal medicine residency programs have problem residents. The presenting characteristics and underlying issues are diverse, so identifying and understanding problem residents is a challenging task. An individualized approach to managing each problem resident may be most successful.

**Acknowledgment:** We are indebted to David Kern, MD, MPH; L. Randol Barker, MD, ScM; and Nicholas Fiebach, MD, for their comments on the manuscript and Ken Kolodner, PhD, for his assistance with data analysis.

## REFERENCES

1. American Board of Internal Medicine. In: Materials from Association of Program Directors in Internal Medicine (APDIM)'s Chief Residents' Workshop on Problem Residents; April 19, 1999; New Orleans, La.
2. Shapiro J, Prislun MD, Larsen KM, Lenahan PM. Working with the resident in difficulty. *Fam Med.* 1987; 19:368-375.
3. Steinert Y, Levitt C. Working with the "problem" resident: guidelines for definition and intervention. *Fam Med.* 1993;25:627-632.
4. Gordon MJ. A prerogatives-based model for assessing and managing the resident in difficulty. *Fam Med.* 1993;25:637-645.
5. Samkoff JS, McDermott RW. Emotional impairment in resident physicians. *Pa Med.* 1989;92:40-43.
6. Smith JW, Denny WF, Witzke DB. Emotional impairment in internal medicine house staff. *JAMA.* 1986; 255:1155-1158.
7. Valko RJ, Clayton PJ. Depression in the internship. *Dis Nerv Syst.* 1975;36:26-29.
8. Ford CV. Emotional distress in internship and residency: a questionnaire study. *Psychiatr Med.* 1983; 1:143-150.
9. Reuben DB. Depressive symptoms in medical house officers. *Arch Intern Med.* 1985;145:286-288.

10. Hsu K, Marshall V. Prevalence of depression and distress in a large sample of Canadian residents, interns, and fellows. *Am J Psychiatry*. 1987;144:1561-1566.
11. McCue JD. The effects of stress on physicians and their medical practice. *N Engl J Med*. 1982;306:458-463.
12. Resident Services Committee, Association of Program Directors in Internal Medicine. Stress and impairment during residency training: strategies for reduction, identification, and management. *Ann Intern Med*. 1988;109:154-161.
13. Landau C, Hall S, Wartman SA, Macko MB. Stress in social and family relationships during the medical residency. *J Med Educ*. 1986;61:654-660.
14. Hurwitz TA, Beiser M, Nichol H, Patrick L, Kozak J. Impaired interns and residents. *Can J Psychiatry*. 1987;32:165-169.
15. Butterfield PS. The stress of residency. *Arch Intern Med*. 1988;148:1428-1435.
16. Quill TE, Williamson PR. Healthy approaches to physician stress. *Arch Intern Med*. 1990;150:1857-1861.
17. O'Connor PG, Spickard AJ. Physician impairment by substance abuse. *Med Clin North Am*. 1997;81:1037-1052.
18. Benzer DG. Managing substance use disorders in resident physicians. *J Am Osteopath Assoc*. 1993;93:217-220.
19. Herrington RE. The impaired physician—recognition, diagnosis, and treatment. *Wis Med J*. 1979;78:21-23.
20. McNamara RM, Margulies JL. Chemical dependency in emergency medicine residency programs: perspective of the program directors. *Ann Emerg Med*. 1994;23:1072-1076.
21. American Medical Association Council on Mental Health. The sick physician: impairment by psychiatric disorders, including alcoholism and drug dependence. *JAMA*. 1973;223:684-687.
22. Aach RD, Girard DE, Humphrey H, et al. Alcohol and other substance abuse and impairment among physicians in residency training. *Ann Intern Med*. 1992;116:245-254.
23. Hughes PH, Conard SE, Baldwin DC Jr, et al. Resident physician substance use in the United States. *JAMA*. 1991;265:2069-2073.
24. Jex SM, Hughes P, Storr C, et al. Relations among stressors, strains, and substance use among resident physicians. *Int J Addict*. 1992;27:979-994.
25. Reuben DB, Noble S. House officer responses to impaired physicians. *JAMA*. 1990;263: 958-960.
26. *Graduate Medical Education Directory*. Chicago, Ill: American Medical Association; 2000.
27. *STATA 6.0* [computer program]. College Station, Tex: Stata Corp; 1999.
28. *The Problem Resident* [videotape]. Philadelphia, Pa: American Board of Internal Medicine; 1992.
29. Boote RM. Guidelines for training programs for staying out of legal trouble: a systematic approach to the problem resident. Workshop presented at: Society of General Internal Medicine national meeting; April 29, 1999; San Francisco, Calif.
30. American Board of Internal Medicine. *Project Professionalism*. Philadelphia, Pa: American Board of Internal Medicine; 1995.
31. vanIneveld CH, Cook DJ, Kane SL, King D. Discrimination and abuse in internal medicine residency. *J Gen Intern Med*. 1996;11:401-405.
32. Balon R, Mufti R, Williams M, Riba M. Possible discrimination in recruitment of psychiatry residents? *Am J Psychiatry*. 1997;154:1608-1609.

That our planet is the one and only planet where life has emerged would be a ridiculous assumption . . . Even if only one in a hundred of the ten billion suitable planets has actually got life well under way, there would be more than 100 million such planets. No, it is presumptuous to think that we are alone.  
 —Harlow Shapley (1885-1972)